

## QUADRATIC THEORY IN BRIEF.

$$\text{Given } ax^2 + bx + c = 0$$
$$\text{then } x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

1. Use the quadratic formula to solve these equations (solutions to 2 dec pl.)

(a)  $3x^2 + 9x + 5 = 0$   
 $ax^2 + bx + c = 0$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$
$$= \frac{-9 \pm \sqrt{9^2 - 4 \times 3 \times 5}}{2 \times 3}$$
$$= \frac{-9 \pm \sqrt{21}}{6}$$

$$x = -0.74 \text{ or } -2.26$$

(b)  $5x^2 - 7x - 11 = 0$   
 $ax^2 + bx + c = 0$        $a = 5 \text{ but } b = -7 \text{ and } c = -11$


$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$
$$= \frac{+7 \pm \sqrt{49 - 4 \times 5 \times (-11)}}{2 \times 5}$$
$$= \frac{+7 \pm \sqrt{49 - 4 \times 5 \times (-11)}}{2 \times 5}$$
$$= \frac{+7 \pm \sqrt{269}}{10}$$

$$x = 2.34 \text{ or } -0.94$$

---

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

THE DISCRIMINANT       $\Delta = b^2 - 4ac$



**REMEMBER:** The solutions of an equation are where the graph of the equation crosses the x axis.

## “COMPLETING THE SQUARE” METHOD.


2. Show clearly how to solve each of the following 4 equations by completing the square (even though 2 of them factorise) and state how the **discriminant** affects the type of solutions.

$$\begin{aligned}\text{(a)} \quad x^2 - 8x + 7 &= 0 \\ x^2 - 8x &= -7 \\ x^2 - 8x + 16 &= -7 + 16 \\ (x - 4)^2 &= 9\end{aligned}$$

$$\begin{aligned}\text{so } x - 4 &= 3 \text{ or } x - 4 = -3 \\ x &= 7 \text{ or } 1\end{aligned}$$

*The solutions are where the graph crosses the x axis.*


$$\begin{aligned}\Delta &= 8^2 - 4 \times 1 \times 7 \\ &= 64 - 28 \\ &= 36 \text{ (a perfect square) so we get 2 rational sols}\end{aligned}$$




$$\begin{aligned}\text{(b)} \quad x^2 - 8x + 16 &= 0 \\ x^2 - 8x &= -16 \\ x^2 - 8x + 16 &= -16 + 16 \\ (x - 4)^2 &= 0 \\ x &= 4\end{aligned}$$

*The solutions are where the graph crosses the x axis.*

$$\begin{aligned}\Delta &= 8^2 - 4 \times 1 \times 16 \\ &= 64 - 64 \\ &= 0 \text{ so we get 1 rational sol.}\end{aligned}$$



$$\begin{aligned}
 \text{(c)} \quad x^2 - 8x + 5 &= 0 \\
 x^2 - 8x &= -5 \\
 x^2 - 8x + 16 &= -5 + 16 \\
 (x - 4)^2 &= 11 \\
 x - 4 &= \pm\sqrt{11} \\
 x &= 4 \pm \sqrt{11} \\
 &\approx 7.32, 0.683
 \end{aligned}$$




The solutions are where the graph crosses the x axis.

$$\begin{aligned}
 \Delta &= 8^2 - 4 \times 1 \times 5 \\
 &= 64 - 20 \\
 &= 44 \quad \text{so we get 2 irrational sols. because 44 does not have} \\
 &\quad \text{an exact square root.}
 \end{aligned}$$


---

$$\begin{aligned}
 \text{(d)} \quad x^2 - 8x + 20 &= 0 \\
 x^2 - 8x &= -20 \\
 x^2 - 8x + 16 &= -20 + 16 = -4 \\
 (x - 4)^2 &= -4 \\
 &\text{Can't find } \sqrt{-4}
 \end{aligned}$$



The solutions are where the graph crosses the x axis but it does not cross the x axis so there are no real solutions.

$$\Delta = 64 - 80 = -16 \text{ so no real sols.}$$


---

3. The Discriminant is  $\Delta = b^2 - 4ac$ .

State what **type** of solutions you get if the discriminant is :

(a) 0  
= 1 rat sol  
(graph sits on x axis)

(b) 1 or 4 or 9 or 16 etc  
= 2 rat sol  
(graph crosses x axis at whole numbers or fractions)

(c) 2 or 3 or 5 or 6 etc  
= 2 irrat sol  
(graph crosses x axis at numbers which are SURDS (eg  $\sqrt{3}$ ))

(d) -1 or -5 or -76 etc  
= NO real solutions  
(graph does not cross x axis)

## EXAMPLES

1.

Find the **value** of  $p$  so that


$$x^2 - 10x + p = 0 \text{ has one solution.}$$

This will have only 1 solution if the graph sits on the  $x$  axis.

In which case, the discriminant  $= 0$

$$\begin{aligned} \Delta &= 100 - 4p = 0 \\ 100 &= 4p \\ 25 &= p \end{aligned}$$

*Note: if  $p = 25$ , the equation is  $x^2 - 10x + 25 = 0$   
so that  $(x - 5)^2 = 0$   
and the only solution is  $x = 5$*




2.

Find  $p$  so that  $x^2 + (p+2)x + (3p-2) = 0$   
has only one rational solution.

This will have only 1 solution if the graph sits on the  $x$  axis.

In which case, the discriminant  $= 0$

$$\begin{aligned} \Delta &= (p+2)^2 - 4(3p-2) = 0 \\ p^2 + 4p + 4 - 12p + 8 &= 0 \\ p^2 - 8p + 12 &= 0 \\ (p-2)(p-6) &= 0 \\ p &= 2 \text{ or } 6 \end{aligned}$$



Some students find this “double” answer confusing:

It means that if  $p = 2$  the equation  $x^2 + (p+2)x + (3p-2) = 0$   
becomes  $x^2 + 4x + 4 = 0$

and THIS equation only has 1 solution ( $x = -2$ )

AND

It means that if  $p = 6$  the equation  $x^2 + (p+2)x + (3p-2) = 0$   
becomes  $x^2 + 8x + 16 = 0$

and THIS equation only has 1 solution ( $x = -4$ )

